*Highlights on Spanish Astrophysics IX, Proceedings of the XII Scientific Meeting of the Spanish Astronomical Society held on July 18 – 22, 2016, in Bilbao, Spain. F. Figueras, A. Sánchez-Lavega, S. Pérez-Hoyos, A. Alonso Herrero, S. Arribas, C. Hernández Monteagudo (eds.)*

arXiv:1610.03320v1 [astro-ph.SR] 7 Oct 2016

# The Galactic O-Star Catalog (GOSC) and the Galactic O-Star Spectroscopic Survey (GOSSS): current status

**Jesús Maíz Apellániz[1], Álvaro Alonso Moragón[2], Lucía Ortiz de Zárate Alcarazo[2], and the GOSSS team**

[1] Centro de Astrobiología, CSIC-INTA, Madrid, Spain
[2] Universidad Complutense, Madrid, Spain

## Abstract

We present the updates of the Galactic O-Star Catalog (GOSC) that we have undertaken in the last two years: new spectral types, more objects, additional information, and coordination with CDS. We also present updates for the Galactic O-Star Spectroscopic Survey (GOSSS). A new paper (GOSSS-III) has been published and ∼1000 targets have been observed since 2014. Four new setups have been added to our lineup and for two of them we have already obtained over 100 spectra: with OSIRIS at the 10.4 m GTC we are observing northern dim stars and with FRODOspec at the 2.0 m Liverpool Telescope we are observing northern bright stars. Finally, we also make available new versions of MGB, the spectral classification tool associated with the project, and of the GOSSS grid of spectroscopic standards.

## 1 What are GOSC and GOSSS?

GOSC, the Galactic O-Star Catalog, is an on-line catalog for O stars (and relatives): `http://gosc.iaa.es`. The first version was presented by Maíz Apellániz et al. (2004) and the first major update by Sota et al. (2008). The current version is v3.2.2 an it has two sections: one public with 601 objects and another private with 7000+ objects.

GOSSS, the Galactic O-Star Spectroscopic Survey (Maíz Apellániz et al. 2011), is observing all optically accessible Galactic stars that anybody has ever classified as O (if we get time on a large enough telescope). It obtains $R \sim 2500$ spectroscopy in the blue-violet region with S/N $\sim$ 250. It has currently observed 2500+ stars, including all O stars down to complete to $B = 8$, with some objects as dim as $B = 17$. In some cases, mostly for extreme SB2s and variables, we have multiple epochs. As of 2016 we have published three

major blocks of the survey: GOSSS-I (Sota et al. 2011), GOSSS-II (Sota et al. 2014), and GOSSS-III (Maíz Apellániz et al. 2016).

## 2 GOSC+GOSSS goals

The primary goal of GOSC and GOSSS is the spectral classification of O stars. More specifically we aim to [a] identify and classify all optically accessible Galactic O stars, [b] improve classification criteria and possibly define new special types and [c] identify objects wrongly classified as O. They also have five secondary goals:

1. Derive the physical properties of O stars such as effective temperature or $v \sin i$.
2. Study SB2s in collaboration with their high-resolution sister surveys (OWN, CAFÉ-BEANS, IACOB, and NoMaDS).
3. Improve our knowledge of the extinction law and the ISM.
4. Measure the spatial distribution of massive stars and dust.
5. Obtain the massive-star IMF.

## 3 What is new in GOSC in the last two years?

In the last two years GOSC has produced several new versions, going from from v3.1.1 to v3.2.2. The most important update has been the use of GOSSS DR2.0 from GOSSS-III (Maíz Apellániz et al. 2016) for public spectral types (those in the main catalog and supplements 1-4, see Table 1). Furthermore, the following updates have also been included:

- There are 1500 new objects, including all bright B stars.
- We have added WISE H II regions in the clusters and nebulae field.
- Coordinates have been revised with new Tycho-2 and 2MASS data.
- We have added 2MASS catalog numbers and Simbad names.
- We have added identifiers (CPD, BD, and ALS).
- We are coordinating with the CDS team at Strasbourg for updates on Simbad spectral types.

Table 1: Main catalog and supplements in GOSC.

| Name | Public? | Description |
|---|---|---|
| Main | Yes | Galactic O stars |
| S01 | Yes | Galactic WR and WR+O systems |
| S02 | Yes | Other Galactic early-type stars |
| S03 | Yes | Galactic late-type stars |
| S04 | Yes | Extragalactic massive stars |
| S05 | No | Galactic O stars |
| S06 | No | Galactic WR and WR+O systems |
| S07 | No | Other Galactic early-type stars |
| S08 | No | Galactic late-type stars |
| S09 | No | Extragalactic massive stars |
| S10 | No | Galactic early-type candidates not observed with GOSSS |
| S11 | No | Galactic late-type candidates not observed with GOSSS |
| S12 | No | Extragalactic candidates not observed with GOSSS |
| S13 | No | Close companions unresolved in GOSSS |

## 4 What is new in GOSSS in the last two years?

In the last two years the most important addition to GOSSS has been the publication of GOSSS-III (Maíz Apellániz et al. 2016), see section 5. We have also produced new versions of MGB (section 6) and of the OB2500 standard grid (section 7).

We have also observed ∼1000 objects in the last two years alone and we have added four new telescopes to our lineup: GTC, Liverpool, SOAR, and Gemini South. Of those, we emphasize GTC and Liverpool, as they have become two of our workhorses, observing hundreds of objects with each one of them. At the 10.4 m GTC we use OSIRIS to observe northern dim stars ($14 < B < 17$) with filler programs. At the robotic 2.0 m Liverpool we are using FRODOspec, a miniIFU spectrograph, to observe northern bright stars ($11 < B$). For the latter case we have developed a specific pipeline, as all other setups use long slits.

## 5 GOSSS-III

We have recently published the third major block of the GOSSS project (Maíz Apellániz et al. 2016). It includes the second of the project data releases (GOSSS DR2.0). The paper presents 142 additional stellar systems with O stars, for a total of 590 in GOSSS DR2.0. Among them, there are 20 new O stars and 11 new SB2 systems: 6 of O+O type and 5 of O+B type. GOSSS-III also shows revisions of previous GOSSS spectral types, including adaptations to the OV/OVz scheme of Arias et al. (2016). We also present egregious errors in the literature i.e. late-type stars classified as O type and introduce luminosity class IV for spectral types O4-O5.5.




Figure 1: Two examples of GOSSS GTC spectra. [top] A newly discovered O7.5 III((f))p star, whose spectrum likely reveals a strong magnetic field, a rarity among O stars. [bottom] A new O3 V((f*)) + O9.5 V: spectroscopic binary system. The primary is the earliest known O dwarf in the northern hemisphere.




Figure 2: [left] Wavelength-collapsed FRODOspec@Liverpool image of the multiple system HD 5005 A+B+C (a D component is located just outside the field to the right). The letters indicate the positions of the three stars. The field size is 9.″84×9.″84 and corresponds to a grid of 12×12 fibers. [right] Pipeline-extracted spectra of the three stars. The pipeline fits a triple Moffat profile at each wavelength point and iterates after fitting a polynomial in wavelength to the PSF parameters.




Figure 3: Example of fitting an SB2 system with MGB. Eight parameters can be adjusted: the spectral subtypes, luminosity classes, and velocities of both the primary and secondary, the flux fraction of the secondary, and the rotation index n. Here HD 93 161 A (black) is fitted with a combination (red) of 60% of HD 152 590 and 40% of 10 Lac separated by 325 km/s.

## 6 MGB

MGB is a code that attacks spectral classification: Maíz Apellániz et al. (2012). It can be used to apply classical visual (non-automatic) spectral classification by interactively comparing with a standard grid. Four parameters can be adjusted:

- Spectral subtype (horizontal classification).
- Luminosity class (vertical classification).
- n index (broadening).
- Alternative standards at each grid point (e.g. ONC or f variants).

The code includes fitting of SB2 systems (Figure 3).

Here we introduce a new version of the code, MGB v2.0, which is available now from `http://jmaiz.iaa.es`. The latest version includes a new default standard grid, OB2500 v3.0, which covers the O2-O9.7 spectral subtypes with GOSSS data (see below). Other grids (O-type or other) from different on-going high-resolution surveys (e.g. IACOB, OWN, IACOBsweG) or from atmosphere models can also be used.

## 7 The GOSSS standard grid

We present a new version of the GOSSS standard grid, OB2500 v3.0, which is integrated with MGB (see above). It covers the spectral subtypes from O2 to O9.7 and the luminosity classes from Vz to Ia (Table 2). There are two types of gaps in the grid: non-existing types (blank) and standards not yet found (...). It is similar to OB2500 v2.0, the grid in Maíz Apellániz (2015), but with small changes introduced by Maíz Apellániz et al. (2016): addition of the Vz luminosity class, definition of luminosity class IV for O4-O5.5, and introduction of new standards. It is available from `http://jmaiz.iaa.es` with MGB v2.0. We plan to have a future extension to A0 (including all B stars) and luminosity class Ia+.

## 8 The GOSSS future

Our plans for the future include:

1. We will extend GOSSS from 590 O stars to ∼1500 in the next decade.
2. We will publish another 3000 spectra and spectral types (mostly B stars) in the same period.
3. We expect to discover several tens of new SB2 systems.
4. We will combine our results with Gaia and other surveys to derive the 6-D distribution of O stars in the solar neighborhood.
5. We will use photometric surveys (e.g. GALANTE) to expand the sample.
6. A complete sample is expected for the two outer Galactic quadrants (low extinction) at the end of survey. The two inner quadrants require (a) multifiber surveys such as WEAVE and (b) IR surveys.
7. We will study the spatial distribution of dust as a function of grain size.
8. We will produce standard grids for the whole OB spectral range.

## References


- Arias, J. I. et al. 2016, *AJ* **152**, 31.
- Maíz Apellániz, J. et al. 2004, *ApJS* **151**, 103.
- Maíz Apellániz, J. et al. 2011, *Highlights of Spanish Astrophysics VI*, 467.
- Maíz Apellániz, J. et al. 2012, *ASPC* **465**, 484.
- Maíz Apellániz, J. et al. 2015, *Highlights of Spanish Astrophysics VIII*, 603.

- Maíz Apellániz, J. et al. 2016, *ApJS* **224**, 4 (GOSSS-III).
- Sota, A. et al. 2008, *RevMexAA (SC)* **33**, 56.
- Sota, A. et al. 2011, *ApJS* **193**, 24 (GOSSS-I).
- Sota, A. et al. 2014, *ApJS* **211**, 10 (GOSSS-II).

Table 2. The OB2500 v3.0 grid of standards.

| | Vz | V | IV | III | II | Ib | Iab/I | Ia |
|---|---|---|---|---|---|---|---|---|
| O2 | | | | | | | *HD 93 129 AaAb* | |
| O3 | *HD 64 568* | ··· | | ··· | | | Cyg OB2-7 | |
| O3.5 | *HD 93 128* | ··· | | *Pismis 24-17* | | | *NGC 3603 HST-48* | |
| O4 | *HD 96 715* | **HD 46 223** | **HD 168 076 AB**<br>*HD 93 250 AB* | ··· | | | HD 15 570<br>HD 16 691<br>HD 190 429 A | |
| O4.5 | ··· | HD 15 629<br>*HDE 303 308 AB* | HD 193 682 | ··· | | | HD 14 947<br>Cyg OB2-9 | |
| O5 | **HD 46 150** | *HDE 319 699* | **HD 168 112 AB** | *HD 93 843* | | | *CPD -47 2963 AB* | |
| O5.5 | ··· | *HD 93 204* | ··· | ··· | | | Cyg OB2-11<br>*ALS 18 747* | |
| O6 | HD 42 088<br>*HDE 303 311* | **ALS 4880**<br>*CPD -59 2600* | *HD 101 190 AaAb* | HDE 338 931 | HDE 229 196 | ··· | ··· | **HD 169 582** |
| O6.5 | *HD 91 572* | **HD 167 633**<br>HD 12 993 | *HDE 322 417* | HD 190 864<br>*HD 96 946*<br>*HD 152 723 AaAb*<br>*HD 156 738 AB* | **HD 157 857** | ··· | ··· | *HD 163 758* |
| O7 | *HD 97 966*<br>*CPD -58 2620*<br>HDE 242 926<br>*HD 91 824* | *HD 93 146 A*<br>*HD 93 222 AB* | ALS 12 320 | Cyg OB2-4 A<br>*HD 93 160 AB* | *HD 94 963*<br>*HD 151 515* | *HD 69 464*<br>HD 193 514 | ··· | ··· |
| O7.5 | *HD 152 590* | HD 35 619 | *HD 97 319* | *HD 163 800* | HD 34 656<br>**HD 171 589** | HD 17 603<br>*HD 156 154* | HD 192 639<br>**9 Sge** | ··· |
| O8 | *HDE 305 539*<br>*HDE 305 438* | *HD 101 223*<br>*HD 97 848*<br>HD 191 978 | *HD 94 024*<br>*HD 135 591* | *HDE 319 702*<br>**$\lambda$ Ori A** | *63 Oph* | **BD -11 4586** | HD 225 160 | *HD 151 804* |
| O8.5 | | *HDE 298 429*<br>HD 14 633 AaAb<br>**HD 46 149**<br>*HD 57 236*<br>*Trumpler 14-9* | **HD 46 966 AaAb** | *HD 114 737 AB*<br>HD 218 195 A | *HD 75 211*<br>HD 207 198 | *HD 125 241* | ··· | *HDE 303 492* |
| O9 | | 10 Lac<br>HD 216 898<br>*CPD -59 2551* | *HD 93 028*<br>*CPD -41 7733* | *HD 93 249 A*<br>HD 24 431 | *HD 71 304*<br>*$\tau$ CMa AaAb* | 19 Cep | HD 202 124<br>*HD 152 249*<br>HD 210 809 | $\alpha$ Cam |
| O9.2 | | **HD 46 202**<br>HD 12 323 | *HD 96 622* | *CPD -35 2105 AaAbB*<br>HD 16 832 | ALS 11 761 | *HD 76 968* | *HD 154 368*<br>*HD 123 008*<br>HD 218 915 | *HD 152 424* |
| O9.5 | | AE Aur<br>*$\mu$ Col* | HD 192 001<br>*HD 93 027*<br>*HD 155 889 AB* | *HD 96 264* | **$\delta$ Ori AaAb** | ··· | HD 188 209 | ··· |
| O9.7 | | *$\upsilon$* **Ori** | HD 207 538 | HD 189 957<br>*HD 154 643* | *HD 68 450*<br>*HD 152 405*<br>HD 10 125 | **HD 47 432**<br>*HD 154 811*<br>*HD 152 147* | HD 225 146<br>*$\mu$ Nor*<br>*HD 104 565*<br>HD 191 781 | HD 195 592<br>*GS Mus* |

Notes Normal, *italic*, and **bold** typefaces are used for stars with $\delta > +20°$, $\delta < -20°$, and the equatorial intermediate region, respectively.